

Application of Asymptotic Iteration Method (AIM) to a Deformed Well Problem

Hakan Ciftci^{*a} and H.F. Kisoglu^{†b}

^a*Gazi Üniversitesi, Fen Fakültesi, Fizik Bölümü, 06500 Teknikokullar Ankara, Türkiye*

^b*Physics Department, Faculty of Science and Letters, Aksaray University, Aksaray, Turkey*

June 24, 2015

Abstract

We have used Asymptotic Iteration Method (AIM) for obtaining the eigenvalues of the Schrödinger's equation for a deformed well problem representing trigonometric functions. By solving the problem, we have found that the Schrodinger's equation for the considered potential has quasi-exact solutions. Additionally, we have also calculated the perturbation expansion of energy eigenvalues and found very simple analytical expression of the energy. Finally, we have considered more general cases and obtained energy eigenvalues for arbitrary potential parameters.

Key Words: Asymptotic iteration method, quasi-exact solution, perturbation method, approximate solution

1 Introduction

The solution method of the Schrödinger's equation, which is the main scope of quantum mechanics, depends upon the configuration of the system held, i.e., behavior of the potential. One of the advantages of the asymptotic iteration method (AIM), introduced in [1, 13], is that it can be used to solve the Schrödinger's equation exactly [2, 6, 7, 14–16] or approximetly for numerous potentials [3, 4, 8, 10, 11, 17].

AIM can also be used in the framework of perturbation theory. In common perturbation theory, eigenfunctions of unperturbed hamiltonian (i.e. exact solvable part) is required to construct the perturbation expansion for energy eigenvalues and eigenfunctions. Furthermore, it is big challenging even for third or fourth-order of approximation in some cases. The advantage of the perturbation method using AIM is that the coefficients in perturbation expansion can be calculated directly without any need to eigenfunctions of unperturbed hamiltonian [5, 9–12].

We use the AIM to find out both the quasi-exact and approximate (numeric) solutions of the Schrödinger's equation for a deformed well problem representing trigonometric functions in the present work. The AIM is summarized in section II. We find out the

^{*}hciftci@gazi.edu.tr

[†]hasanfatihk@aksaray.edu.tr

eigenvalues of Schrödinger's equation approximately and quasi-exactly in section III for a potential defined by

$$V(x) = A \cos(x) + B^2 \sin^2(x) + \frac{\gamma(\gamma+1)}{\sin^2(x)}, \quad x \in (0, \pi) \quad (1)$$

where A and B are coefficients and γ ranges over $(0, \infty)$. In section IV, we apply perturbation method for this kind of potential, just after, summarizing the perturbation theory in the framework of AIM.

2 The Asymptotic Iteration Method (AIM)

Although the asymptotic iteration method has been introduced comprehensively in [1], we outline this method in this section according to organization of the paper. The AIM can be used to solve second-order homogeneous linear differential equations of the form

$$y''(x) = \lambda_0(x)y'(x) + s_0(x)y(x) \quad (2)$$

where λ_0 and s_0 have continuous derivatives in the defined interval of the x independent variable. According to asymptotic aspect of the method, if we have

$$\frac{s_n}{s_{n-1}} = \frac{\lambda_n}{\lambda_{n-1}} \equiv \alpha \quad (3)$$

for sufficient large $n > 0$. The general solution of Eq.2 is obtained as

$$y(x) = \exp\left(-\int^x \alpha(t)dt\right) \left[C_2 + C_1 \int^x \exp\left(\int^t (\lambda_0(t) + 2\alpha(t)) d\tau\right) dt \right] \quad (4)$$

where

$$\begin{aligned} \lambda_n &= \lambda'_{n-1} + s_{n-1} + \lambda_0 \lambda_{n-1} \\ s_n &= s'_{n-1} + s_0 \lambda_{n-1} \end{aligned} \quad (5)$$

Eigenvalues, E , of the Schrödinger's equation are obtained using the following termination condition by means of Eq.3

$$\delta_n(x) = s_n(x)\lambda_{n-1}(x) - \lambda_n(x)s_{n-1}(x) = 0 \quad (6)$$

where the eigenvalues of n energy levels are obtained by Eq.6 with n iterations, if the problem is analytically solvable (exact solution). In this case, eigenvalues and eigenfunctions can be determined in explicit algebraic form in AIM. Furthermore, there are

limited number of suitable potentials for this case.

In exact solvable problem, condition $\delta_n = 0$ is satisfied at every x point in the defined interval since δ_n is independent from x . The (unknown) E values are determined from the n roots of condition $\delta_n = 0$. In case of approximation, δ_n depends on both x and E . So, we have to determine a $x = x_0$ initial value for solving $\delta_n = 0$ with respect to E . As this initial x_0 value may be found from the minimum value of the potential may also bounded domain of the eigenvalue problem [3, 11].

3 AIM for the Problem

We have considered a deformed well problem representing trigonometric functions as below

$$V(x) = A \cos(x) + B^2 \sin^2(x) + \frac{\gamma(\gamma+1)}{\sin^2 x}, \quad x \in (0, \pi)$$

where A , B and γ are the potential parameters. γ is defined in the interval $[0, \infty)$. If we substitute $V(x)$ in the Schrödinger's equation, we have

$$\left\{ \frac{d^2}{dx^2} + E - \left[A \cos(x) + B^2 \sin^2(x) + \frac{\gamma(\gamma+1)}{\sin^2(x)} \right] \right\} \Psi(x) = 0 \quad (7)$$

where unnormalized eigenfunction $\Psi(x) \in L^2(0, \pi)$. We may assume that $\Psi(x)$ takes the form

$$\Psi(x) = \sin^{\gamma+1}(x) e^{B \cos(x)} f(x) \quad (8)$$

with this choice, $\Psi(x)$ vanishes at $x = 0$ and $x = \pi$. Once we substitute above equation in Eq.7

$$f''(x) + 2 \left[(\gamma+1) \frac{\cos(x)}{\sin(x)} - B \sin(x) \right] f'(x) + \left[E - (\gamma+1)^2 - \{A + B(2\gamma+3)\} \cos(x) \right] f(x) = 0 \quad (9)$$

is obtained. If we change the variable such as $\cos(x) = y$ in Eq.9 which yields

$$f''(y) = 2 \left(\frac{\sigma y}{1-y^2} - B \right) f'(y) + \left(\frac{\xi y - \omega}{1-y^2} \right) f(y) \quad (10)$$

where $\omega = E - (\gamma+1)^2$, $\xi = A + 2\sigma B$ and $2\sigma = 2\gamma+3$. The last equation is in form of Eq.2 for which AIM can be used. We obtain the (unkown) E energy eigenvalues by using AIM.

3.1 Quasi-Exact Solutions of the Problem

We have obtained Eq.10 for using the AIM for determining the E eigenvalues of Eq.7. Our calculations show that the problem has quasi-exact solution under some special

conditions on the potential parameters. AIM gives us that if $\xi = -2nB$ with $n = 0, 1, 2, 3, \dots$ the energy eigenvalues can be obtained exactly. We will give first two results analytically because of the complecity of the calculations, but we will give more numerical results in Table.1. Using $\xi = -2nB$ and $\xi = A + 2\sigma B$ together, we get the following formula

$$A = -2B \left(n + \gamma + \frac{5}{2} \right) \quad (11)$$

For $n = 0$, we get $\omega = 0$ and $E_{00} = (\gamma + 1)^2$. For $n = 1$, we get $\omega = \sigma \mp \sqrt{\sigma^2 + 4B^2}$ and the energies can be obtained as follows

$$E_{10} = \left(\gamma + \frac{3}{2} \right)^2 + \frac{1}{4} - \sqrt{\left(\gamma + \frac{3}{2} \right)^2 + 4B^2} \quad (12)$$

$$E_{11} = \left(\gamma + \frac{3}{2} \right)^2 + \frac{1}{4} + \sqrt{\left(\gamma + \frac{3}{2} \right)^2 + 4B^2} \quad (13)$$

Numerically calculated eigenvalues for $B = 1$ and $\sigma = 4.5$ can be found in Table.1.

Table 1. Numerically calculated eigenvalues for $B = 1$ and $\sigma = 4.5$ for $n = 0$ to 5 by using $\xi = -2nB$. k is sub-level for each n level and $k=0, 1, 2, \dots, n$.

n	k	A	ω_{nk}	E_{nk}
0	0	-11	0	16.0000
1	0	-13	-0.424429	15.5756
	1	-13	9.42443	25.4244
2	0	-15	-0.919071	15.0809
	1	-15	9.27711	25.2771
	2	-15	20.642	36.642
3	0	-17	-1.47815	14.5218
	1	-17	9.09552	25.0955
	2	-17	20.6205	36.6205
	3	-17	33.7621	49.7621
4	0	-19	-2.09614	13.9039
	1	-19	8.87768	24.8777
	2	-19	20.5892	36.5892
	3	-19	33.7953	49.7953
	4	-19	48.834	64.834
5	0	-21	-2.76786	13.2321
	1	-21	8.62229	24.6223
	2	-21	20.5459	36.5459
	3	-21	33.8293	49.8293
	4	-21	48.8905	64.8905
	5	-21	65.8799	81.8799

3.2 Approximately Determination of the Eigenvalues

In this case, we have determined the eigenvalues for arbitrary A , B and γ values. Calculated eigenvalues (E_{direct}) of $n = 0$ to 5 are given in Table.2 in which A , B and γ (in

$(A; B; \gamma)$ form) are $(0.167; 0.0019; 0.0019)$, $(2; 0.5; 1)$ and $(4; 1; 1.97)$. In the calculation we take $y_0 = \frac{1}{2}$.

Table 2. Approximately calculated eigenvalues E_{direct} for $(0.167; 0.0019; 0.0019)$, $(2; 0.5; 1)$ and $(4; 1; 1.97)$ in $(A; B; \gamma)$ form. The value written as subscript for each eigenvalue represents the iteration number.

	$A = 0.167, B = \gamma = 0.0019$		$A = 2, B = 0.5, \gamma = 1$		$A = 4, B = 1, \gamma = 1.97$	
n	ω	E_{direct}	ω	E_{direct}	ω	E_{direct}
0	-0.00231	1.00149 ₍₃₎	0.07399	4.07399 ₍₁₀₎	0.57328	9.39418 ₍₁₃₎
1	3.00473	4.00853 ₍₄₎	5.17104	9.17104 ₍₁₂₎	7.618	16.4389 ₍₁₃₎
2	8.008	9.0118 ₍₅₎	12.1609	16.1609 ₍₁₁₎	16.5264	25.3473 ₍₁₄₎
3	15.0116	16.0154 ₍₆₎	21.1506	25.1506 ₍₁₃₎	27.4331	36.254 ₍₁₆₎
4	24.0153	25.0191 ₍₇₎	32.1438	36.1438 ₍₁₄₎	40.348	49.1689 ₍₁₈₎
5	35.0191	36.0229 ₍₈₎	45.1392	49.1392 ₍₁₅₎	55.2698	64.0907 ₍₁₉₎

As is seen from Table.2, energy eigenvalues act as $\sim (n + \gamma + 1)^2$.

4 Perturbation Theory within the frame of AIM

Although the usage of perturbation method within AIM is introduced in [9] comprehensively, we give a summary about the application method in this section. Assume that the potential of a system is written in a form of

$$V(x) = V_0(x) + \mu V_p(x) \quad (14)$$

where $V_0(x)$ is solvable (or unperturbed hamiltonian) potential. $V_p(x)$ and μ are potential of the perturbed hamiltonian and perturbation expansion parameter, respectively. So, the Schrödinger's equation reads

$$\left(-\frac{d^2}{dx^2} + V_0(x) + \mu V_p(x) \right) \Psi(x) = E \Psi(x) \quad (15)$$

where E_n eigenvalues are written as the expansion of j th-order correction $E_n^{(j)}$ as follows

$$E_n = E_n^{(0)} + \mu E_n^{(1)} + \mu^2 E_n^{(2)} + \dots = \sum_{j=0}^{\infty} \mu^j E_n^{(j)} \quad (16)$$

The form of Eq.2 can be yielded by a substitution of (in general form) $\psi(x) = \psi_0(x)f(x)$ in Eq.15. So, one can obtain the following equation for $f(x)$

$$f''(x) = \lambda_0(x, \mu, E) f'(x) + s_0(x, \mu, E) f(x) \quad (17)$$

The termination condition, Eq.6, is written as

$$\delta_n(x, \mu, E) = s_n(x, \mu, E) \lambda_{n-1}(x, \mu, E) - \lambda_n(x, \mu, E) s_{n-1}(x, \mu, E) = 0 \quad (18)$$

Once $\delta_n(x, \mu, E)$ is expanded about $\mu = 0$, we obtain

$$\delta_n(x, \mu, E) = \delta_n(x, 0, E) + \frac{\mu}{1!} \frac{\partial \delta_n(x, \mu, E)}{\partial \mu} \Big|_{\mu=0} + \frac{\mu^2}{2!} \frac{\partial^2 \delta_n(x, \mu, E)}{\partial \mu^2} \Big|_{\mu=0} + \dots = \sum_{k=0}^{\infty} \mu^k \delta_n^{(k)}(x, E) = 0 \quad (19)$$

where $\delta_n^{(k)}(x, E) = \frac{1}{k!} \frac{\partial^k \delta_n(x, \mu, E)}{\partial \mu^k} \Big|_{\mu=0}$. The condition of

$$\delta_n^{(k)}(x, E) = 0 \quad (20)$$

can be obtained by means of Eq.19 for each k . According to the perturbation method in the framework of AIM, solving the equation $\delta_n(x, 0, E) = 0$ with respect to (unknown) E gives $E_n^{(0)}$ (eigenvalues of unperturbed hamiltonian), equation $\delta_n^{(1)}(x, E) = 0$ gives $E_n^{(1)}$ (first-order correction to E_n), $\delta_n^{(2)}(x, E)$ gives $E_n^{(2)}$ (second-order correction to E_n) and so on. The eigenfunctions can also be found with similar manner. This is an attractive feature of the AIM usage in the perturbation theory for obtaining the eigenfunctions $f_n(x)$ given as follows

$$f_n(x) = \exp \left(- \int^x \alpha_n(t, \mu) dt \right) \quad (21)$$

where $\alpha_n(t, \mu) \equiv s_n(t, \mu)/\lambda_n(t, \mu)$. $\alpha_n(t, \mu)$ is expanded about $\mu = 0$ in a similar manner of obtaining the eigenvalues. So,

$$\alpha_n(t, \mu) = \sum_{k=0}^{\infty} \mu^k \alpha_n^{(k)}(t) \quad (22)$$

where $\alpha_n^{(k)}(t) = \frac{1}{k!} \frac{\partial^k \alpha_n(t, \mu)}{\partial \mu^k} \Big|_{\mu=0}$. Thus, perturbation expansion of the $f_n(x)$ is written as follows

$$f_n(x) = \exp \left[\sum_{k=0}^{\infty} \mu^k \left(- \int^x \alpha_n^{(k)}(t) dt \right) \right] = \prod_{k=0}^{\infty} f_n^{(k)}(x) \quad (23)$$

where k th-order correction $f_n^{(k)}(x)$ to $f_n(x)$ is

$$f_n^{(k)}(x) = \mu^k \left(- \int^x \alpha_n^{(k)}(t) dt \right) \quad (24)$$

4.1 Perturbation Theory for the Potential Function in the frame of AIM

In section III, we have obtained the eigenvalue equation in the appropriate form for AIM usage as follows

$$f''(y) = 2 \left(\frac{\sigma y}{1-y^2} - B \right) f'(y) + \left(\frac{\xi y - \omega}{1-y^2} \right) f(y) \quad (25)$$

where $\omega = E - (\gamma + 1)^2$, $\xi = A + 2\sigma B$ and $2\sigma = 2\gamma + 3$. A and B parameters of the potential $V(x)$, given by Eq.1, was linked to eachother via Eq.11. Suppose that $A = 2aB$, in this case $\xi = 2(a + \sigma)B$ and the differential equation can be written as below

$$f''(y) = 2 \left(\frac{\sigma y}{1-y^2} - B \right) f'(y) + \left(\frac{2(a + \sigma)By - \omega}{1-y^2} \right) f(y) \quad (26)$$

Now, B can be taken as perturbation expansion parameter. We can expand ω as below

$$\omega_n = \omega_n^{(0)} + B\omega_n^{(1)} + B^2\omega_n^{(2)} + \dots \quad (27)$$

In this expansion the general form of zeroth-order correction $\omega_n^{(0)}$ is obtained via,

$$\delta^{(0)}(x, 0, \omega_n^{(0)}) = 0 \quad (28)$$

as mentioned previously. The obtained few values of $\omega_n^{(0)}$ are 0 , 2σ , $2 + 4\sigma$, $6 + 6\sigma$, $12 + 8\sigma$, $20 + 10\sigma$, respectively. One can easily write general formula for $\omega_n^{(0)}$ as below

$$\omega_n^{(0)} = n(n-1) + 2n\sigma \quad (29)$$

The first-order correction to ω_n is generalized in the similar manner, and it is found that $\omega_n^{(1)} = 0$ for each n level via the equation $\delta^{(1)}(x, 0, \omega_n^{(1)}) = 0$. The second-order correction to ω_n is generalized as

$$\omega_n^{(2)} = -2 \left(\frac{a^2 g_1(n, \sigma) - g_2(n, \sigma)}{g_3(n, \sigma)} \right) \quad (30)$$

where

$$\begin{aligned} g_1(n, \sigma) &= 2\sigma^2 - (2n + 5)\sigma - (n^2 - n - 3) \\ g_2(n, \sigma) &= 2\sigma^4 + (6n - 5)\sigma^3 + (7n^2 - 11n + 3)\sigma^2 + 4n(n-1)^2\sigma + n^2(n-1)^2 \\ g_3(n, \sigma) &= (2\sigma + 2n + 1)(2\sigma + 2n - 3)(\sigma + n)(\sigma + n - 1) \end{aligned} \quad (31)$$

If Eq.29 and Eq.30 are substituted in Eq.27 one can obtain

$$\omega_n = n(n-1) + 2n\sigma - \left(\frac{A^2 g_1(n, \sigma) - 4B^2 g_2(n, \sigma)}{2g_3(n, \sigma)} \right) \quad (32)$$

where we use $a = \frac{A}{2B}$ substituted in Eq.30 as well. From $\omega_n = E_n - (\gamma + 1)^2$, we get E_n^p (perturbed energy) approximately as follows

$$E_n^p \approx (\gamma + n + 1)^2 - \left(\frac{A^2 g_1(n, \sigma) - 4B^2 g_2(n, \sigma)}{2g_3(n, \sigma)} \right) \quad (33)$$

Numeric results are given in Table.3 in which E_n^p , given in Eq.33, was used for $n = 0$ to 5, and in Table.4 for $n = 0$ to 10. The comparison of E_n^p with numerically calculated eigenvalues E_{direct} for arbitrary A , B and γ is also given in Table.3 and Table.4. The values written as subscript represent the iteration numbers in both tables.

Table 3. Comparison of eigenvalues of perturbed hamiltonian E_n^p with numerically ones E_{direct} for $A = 2$, $B = 2$ and $\gamma = 4$ for $n = 0$ to 5.

n	ω_n	E_{direct}	E_n^p
0	3.6259	28.6259 ₍₁₃₎	28.6364
1	14.1219	39.1219 ₍₁₄₎	39.1329
2	26.8245	51.8245 ₍₁₇₎	51.8308
3	41.6323	66.6323 ₍₁₈₎	66.6353
4	58.5005	83.5005 ₍₁₉₎	83.5015
5	77.406	102.406 ₍₁₉₎	102.406

Table 4. Comparison of the eigenvalues of perturbed hamiltonian E_n^p with numerically ones E_{direct} for $A = 2$, $B = 1$ and $\gamma = 0.0019$ for $n = 0$ to 10.

n	ω_n	E_{direct}	E_n^p
0	0.38637	1.39017 ₍₁₂₎	1.42145
1	3.65921	4.66301 ₍₁₃₎	4.64073
2	8.56603	9.56983 ₍₁₆₎	9.56855
3	15.5445	16.5483 ₍₁₅₎	16.547
4	24.5364	25.5402 ₍₁₆₎	25.5392
5	35.5338	36.5376 ₍₁₈₎	36.5368
6	48.5336	49.5374 ₍₁₈₎	49.5369
7	63.5349	64.5387 ₍₁₉₎	64.5383
8	80.537	81.5408 ₍₂₀₎	81.5404
9	99.5392	100.543 ₍₂₀₎	100.543
10	120.542	121.546 ₍₂₁₎	121.546

It can be seen from Table.3 that numerically calculated eigenvalues E_{direct} are consistent with the ones of perturbed hamiltonian E_n^p . Also, they are more consistent as the energy level n increases, and the values of E_{direct} converge more slowly with increasing n . All these comments apply to Table.4 as well. Furthermore, the eigenvalues act as $\sim (n+\gamma+1)^2$

while $\omega_n \approx n(n+2)$ in Table.4.

In Table.5, values of E_{direct} and E_n^p are inconsonant by selecting $A = B = 5$ and $\gamma = 0.5$.

Table 5. Comparison of the eigenvalues of perturbed hamiltonian E_n^p with numerically calculated ones E_{direct} for $A = B = 5$ and $\gamma = 0.5$ for $n = 0$ to 5. The values written as subscript represent the iteration numbers.

n	ω_n	E_{direct}	E_n^p
0	11.8935	14.1435 ₍₃₁₎	21.0000
1	19.3048	21.5548 ₍₃₄₎	20.8333
2	25.0743	27.3243 ₍₃₆₎	25.7917
3	32.0111	34.2611 ₍₄₀₎	33.375
4	41.5015	43.7515 ₍₃₇₎	43.1667
5	53.2222	55.4722 ₍₃₉₎	55.0476

5 Conclusion

We have used AIM for solving Schrödinger's equation for a deformed well problem representing trigonometric function. In this study, we have applied AIM to quasi-exact solvable eigenvalue problem for obtaining the energy eigenvalues and seen that each energy level n has a sub-level k that $k = 0, 1, 2, \dots, n$. Method has also been used to obtain the eigenvalues approximately by using arbitrary potential parameters. According to results, we can say that the energy eigenvalues behave as $\sim (n + \gamma + 1)^2$ for larger quantum number n , as is seen in Eq.33. AIM can also be used in the framework of perturbation theory. In common perturbation theory, eigenfunctions of unperturbed hamiltonian is required to construct the perturbation expansion for energy eigenvalues and eigenfunctions. Also, determining the third or fourth-order correction to energy eigenvalues is usually quite big challenging. The advantage of the perturbation method using AIM is that the coefficients in perturbation expansion can be calculated directly without any need to eigenfunctions of unperturbed hamiltonian and any complication.

Perturbation method has also been applied to potential function held in scope of the AIM, in this study. According to results, eigenvalues of perturbed hamiltonian are consistent with those of approximately calculated ones for which arbitrary parameters of the potential have been used. However, they are inconsonant for some set of the parameters. Although this inconsistency is true for $n = 0$ to 5, it tends to decrease as n increases. Consequently, it can be said that we can make the eigenvalues tend to specific behavior by selecting appropriate potential parameters.

References

- [1] H. Ciftci, R. L. Hall and N. Saad, J. Phys. A: Math. Gen. **36** (2003) 11807.
- [2] A. V. Turbiner, J. Phys. A: Math. Gen. **45** (2012) 025203.
- [3] H. Ciftci, R. L. Hall and N. Saad, Cent. Eur. J. Phys. **11** (2013) 37.
- [4] M. Aygun, O. Bayrak and I. Boztosun, J. Phys. B: At. Mol. Opt. Phys. **40** (2007) 537.

- [5] T. Barakat, J. Phys. A: Math. Gen. **39** (2006) 823.
- [6] O. Ozer, Progress of Theoretical Physics **121** (2009) 437.
- [7] S.-A. Yahiaoui and M. Bentaiba, Acta Physica Polonica B **42** (2011) 1755.
- [8] O. Ozer and G. Levai, Rom. Journ. Phys. **57** (2012) 582.
- [9] H. Ciftci, R. L. Hall and N. Saad, Phys. Lett. A **340** (2005) 388.
- [10] H. Ciftci, R. L. Hall and N. Saad, Phys. Rev. A **72** (2005) 022101.
- [11] B. Champion, R.L. Hall and N. Saad, Int. J. Mod. Phys. A **23** (2008) 1405.
- [12] H. Ciftci, R. L. Hall and N. Saad, J. Phys. A: Math. Gen. **39** (2005) 2338.
- [13] H. Ciftci, R. L. Hall and N. Saad, J. Phys. A: Math. Gen. **38** (2005) 1147.
- [14] Richard L. Hall, Nasser Saad, K. D. Sen and Hakan Ciftci, Phys. Rev. A **80** (2009) 032507.
- [15] Hakan Ciftci, Richard L. Hall, Nasser Saad and Ebubekir Dogu, J. Phys. A: Math. Theor. **43** (2010) 415206.
- [16] O. Bayrak and I. Boztosun, J. Phys. A: Math. Gen. **39** (2006) 6955.
- [17] Babatunde J. Falaye, Majid Hamzavi, Sameer M. Ikhdair, Chinese Physics Letters **30** (2013) 020305.